



Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis

Dena Ettehad, Connor A Emdin, Amit Kiran, Simon G Anderson, Thomas Callender, Jonathan Emberson, John Chalmers, Anthony Rodgers, Kazem Rahimi

Summary

Background The benefits of blood pressure lowering treatment for prevention of cardiovascular disease are well established. However, the extent to which these effects differ by baseline blood pressure, presence of comorbidities, or drug class is less clear. We therefore performed a systematic review and meta-analysis to clarify these differences.

Method For this systematic review and meta-analysis, we searched MEDLINE for large-scale blood pressure lowering trials, published between Jan 1, 1966, and July 7, 2015, and we searched the medical literature to identify trials up to Nov 9, 2015. All randomised controlled trials of blood pressure lowering treatment were eligible for inclusion if they included a minimum of 1000 patient-years of follow-up in each study arm. No trials were excluded because of presence of baseline comorbidities, and trials of antihypertensive drugs for indications other than hypertension were eligible. We extracted summary-level data about study characteristics and the outcomes of major cardiovascular disease events, coronary heart disease, stroke, heart failure, renal failure, and all-cause mortality. We used inverse variance weighted fixed-effects meta-analyses to pool the estimates.

Results We identified 123 studies with 613 815 participants for the tabular meta-analysis. Meta-regression analyses showed relative risk reductions proportional to the magnitude of the blood pressure reductions achieved. Every 10 mm Hg reduction in systolic blood pressure significantly reduced the risk of major cardiovascular disease events (relative risk [RR] 0·80, 95% CI 0·77–0·83), coronary heart disease (0·83, 0·78–0·88), stroke (0·73, 0·68–0·77), and heart failure (0·72, 0·67–0·78), which, in the populations studied, led to a significant 13% reduction in all-cause mortality (0·87, 0·84–0·91). However, the effect on renal failure was not significant (0·95, 0·84–1·07). Similar proportional risk reductions (per 10 mm Hg lower systolic blood pressure) were noted in trials with higher mean baseline systolic blood pressure and trials with lower mean baseline systolic blood pressure (all $p_{\text{trend}} > 0·05$). There was no clear evidence that proportional risk reductions in major cardiovascular disease differed by baseline disease history, except for diabetes and chronic kidney disease, for which smaller, but significant, risk reductions were detected. β blockers were inferior to other drugs for the prevention of major cardiovascular disease events, stroke, and renal failure. Calcium channel blockers were superior to other drugs for the prevention of stroke. For the prevention of heart failure, calcium channel blockers were inferior and diuretics were superior to other drug classes. Risk of bias was judged to be low for 113 trials and unclear for 10 trials. Heterogeneity for outcomes was low to moderate; the I^2 statistic for heterogeneity for major cardiovascular disease events was 41%, for coronary heart disease 25%, for stroke 26%, for heart failure 37%, for renal failure 28%, and for all-cause mortality 35%.

Interpretation Blood pressure lowering significantly reduces vascular risk across various baseline blood pressure levels and comorbidities. Our results provide strong support for lowering blood pressure to systolic blood pressures less than 130 mm Hg and providing blood pressure lowering treatment to individuals with a history of cardiovascular disease, coronary heart disease, stroke, diabetes, heart failure, and chronic kidney disease.

Funding National Institute for Health Research and Oxford Martin School.

Introduction

Elevated blood pressure is the most important risk factor for death and disability worldwide, affecting more than one billion individuals and causing an estimated 9·4 million deaths every year.¹ Prospective cohort studies have reported a continuous log-linear association between blood pressure and vascular events to a blood pressure of 115/75 mm Hg, with no apparent threshold.² This association seems to exist across large and diverse population groups, including men and women, individuals aged 40–89 years, from different ethnicities, with and without established vascular disease.^{2–4} Despite

this robust observational evidence, whether blood pressure lowering treatment reduces the risk of cardiovascular disease in all patient populations remains unclear.

Although the benefits of blood pressure lowering have long been established in randomised trials of patients with substantially raised blood pressures,^{5–8} evidence for the protective effects of pharmacologically-induced blood pressure reduction in individuals with lower blood pressure or with comorbidities, have been less certain.^{9–12} Furthermore, the best approach to reduce blood pressure remains subject to controversy.^{13–16}

Published Online
December 23, 2015
[http://dx.doi.org/10.1016/S0140-6736\(15\)01225-8](http://dx.doi.org/10.1016/S0140-6736(15)01225-8)

See Online/Comment
[http://dx.doi.org/10.1016/S0140-6736\(15\)01344-6](http://dx.doi.org/10.1016/S0140-6736(15)01344-6)

The George Institute for Global Health (D Ettehad MSc, C A Emdin HBSc, A Kiran PhD, S G Anderson PhD, T Callender MB ChB, Prof K Rahimi DM FRCP) and Clinical Trial Service Unit and Epidemiological Studies Unit (J Emberson PhD), University of Oxford, Oxford, UK; Institute of Cardiovascular Sciences, University of Manchester, Manchester, UK (S G Anderson); King's College Hospital NHS Foundation Trust, London, UK (T Callender); and The George Institute for Global Health, University of Sydney, Sydney, NSW, Australia (Prof J Chalmers PhD, Prof A Rodgers PhD)

Correspondence to: Prof Kazem Rahimi, The George Institute for Global Health, University of Oxford, Oxford OX1 3BD, UK
kazem.rahimi@georgeinstitute.ox.ac.uk

Research in context

Evidence before this study

Although the benefits of blood pressure lowering treatment for prevention of cardiovascular disease are well established, the extent to which these effects differ by baseline blood pressure, presence of comorbidities, or drug class is less clear.

Added value of this study

Our study provides a comprehensive systematic review and meta-analysis of all available large-scale blood pressure lowering randomised trials. Our findings show that pharmacological blood pressure lowering results in similar proportional reductions in risk of cardiovascular disease and death to a mean baseline systolic blood pressure of less than 130 mm Hg. Furthermore, we noted that proportional risk reductions are broadly similar among individuals with or without major cardiovascular comorbidities. Finally, our findings emphasise the fact that, despite the general efficacy of commonly prescribed blood pressure lowering drug classes in preventing cardiovascular disease, there are some significant differences among them in the reduction of risk of specific clinical outcomes. For example, calcium channel blockers seem

to be more effective than other classes of drugs for stroke prevention, and diuretics are more effective for prevention of heart failure.

Implications of all the available evidence

Our study has several implications for clinical practice. First, our findings suggest that blood pressure lowering to levels below those recommended in current guidelines (ie, systolic blood pressure of less than 140 mm Hg) will reduce the risk of cardiovascular disease. Second, by showing no evidence for a threshold below which blood pressure lowering ceases to work, the findings call for blood pressure lowering based on an individual's potential net benefit from treatment rather than treatment of the risk factor to a specific target. Third, the broad consistency of the findings across patients with or without prior vascular disease could help to simplify clinical guidelines for use of blood pressure lowering drugs. Fourth, the differences we identified between classes of drugs support more targeted drug use for individuals at high risk of specific outcomes (eg, calcium channel blocker therapy for individuals at high risk of stroke).

See Online for appendix

Recent major guidelines have reversed a trend toward lower blood pressure thresholds and targets, recommending higher targets and threshold for blood pressure lowering than have previous guidelines.^{14,17,18} The SPRINT trial¹² reported the benefits of blood pressure lowering to 120 mm Hg in some high-risk groups of patients. However uncertainty remains as to whether such benefits hold for high-risk individuals excluded from the trial, especially those with diabetes or cerebrovascular disease.¹²

In this systematic review and meta-analysis, we aimed to combine data from all published large-scale blood pressure lowering trials to quantify the effects of blood pressure reduction on cardiovascular outcomes and death across various baseline blood pressure levels, major comorbidities, and different pharmacological interventions.

Methods

Search strategy and selection criteria

The systematic review and tabular meta-analysis adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines for meta-analyses of interventional studies.¹⁹ We searched MEDLINE from Jan 1, 1966, to July 7, 2015, using an existing search strategy¹⁶ with the terms “antihypertensive agents” or “hypertension” or “diuretics”, “thiazide”, or “angiotensin-converting enzyme” or “angiotensin-converting enzyme inhibitors” or “receptors, angiotensin/antagonists & inhibitors” or “tetrazoles” or “calcium channel blockers” or “vasodilator agents” or the names of all blood pressure lowering drugs listed in the British National Formulary as

keywords or text words or the MeSH term “blood pressure/drug effects”. Search terms used in the MEDLINE search are provided in the appendix (pp 39–41). We restricted our search to clinical trials, controlled clinical trials, randomised controlled trials, or meta-analyses. We applied no language restrictions. Reference lists of eligible studies and related meta-analyses were hand searched to identify further relevant studies.

All randomised controlled trials of blood pressure lowering treatment published between Jan 1, 1966, and Nov 9, 2015, were eligible for inclusion. Eligible studies fell into three categories: first, random allocation of participants to a blood pressure lowering drug or placebo; second, random allocation of participants to different blood pressure lowering drugs; and third, random allocation of participants to different blood pressure lowering targets. To minimise the risk of small-study effects,²⁰ all studies were required to have a minimum of 1000 patient-years of follow-up in each study group, the same criterion applied in the Blood Pressure Lowering Treatment Trialists' Collaboration (BPLTTC).²¹ Trials of antihypertensive drugs for indications other than hypertension were eligible. No trials were excluded because of the presence of baseline comorbidities.

Data analysis

Two researchers (DE and CAE) screened all abstracts identified in the initial search and excluded studies in violation of the inclusion criteria. Full-text articles were subsequently reviewed in duplicate and, in cases of disagreement, consensus was achieved through referral




Figure 1: Study selection

to a third reviewer (KR). An electronic data abstraction form was used to record patient and study characteristics, including sample size, treatment comparisons, baseline blood pressure, blood pressure achieved, and mean blood pressure reduction. If not reported, corresponding authors were contacted to obtain data about baseline and achieved blood pressure by use of individually tailored data request forms.

Data were also extracted for major cardiovascular disease events (defined as fatal and non-fatal myocardial infarction, sudden cardiac death, revascularisation, fatal and non-fatal stroke, and fatal and non-fatal heart failure), coronary heart disease (fatal and non-fatal myocardial infarction and sudden cardiac death, excluding silent myocardial infarction), stroke (fatal and non-fatal, excluding transient ischaemic attacks), heart failure (new diagnosis of heart failure, hospital




Figure 2: Meta-regression plot

Plot shows the percentage risk reduction in major cardiovascular events regressed against the difference in achieved systolic blood pressure between study treatment groups.

admission, or death), renal failure (end-stage renal disease resulting in dialysis, transplantation, or death), and all-cause mortality. Information was extracted about the number of patients in each group with the baseline comorbidities of cardiovascular disease, coronary heart disease, cerebrovascular accident (history of stroke or transient ischaemic attack), type 2 diabetes, heart failure, and chronic kidney disease (self-reported chronic kidney disease or creatinine clearance <30 mL/min). The SPRINT trial¹² reported renal failure for patients with chronic kidney disease at baseline; this was recorded for the entire trial.

For all outcomes, the number of events in each group and the summary statistic (either relative risk [RR] or hazard ratio [HR] and 95% CIs) were extracted. We used HRs preferentially to RRs because they incorporate time-to-event and allow for censoring.²² When neither summary statistic was provided, we calculated RRs from the number of events and participants in each treatment group. Conversely, if the total number of events was missing, we estimated it using the summary statistic and its confidence interval.

We used the Cochrane risk of bias tool to assess the methodological quality of the eligible trials.²³ Selection bias (randomisation and allocation concealment), performance bias (blinding of participants and investigators), detection bias (blinding of outcome adjudicators), attrition bias (differential loss to follow-up), and reporting bias (selective outcome reporting) were judged to be of low, unclear, or high risk for each trial. We then judged each trial as a whole to ascertain whether there was low, unclear, or high risk of bias, based on whether the level of bias in each of the defined domains could have led to material biases in the risk estimates.




Figure 3: Standardised effects of a 10 mm Hg reduction in systolic blood pressure
RR=relative risk.

Statistical analysis

We calculated overall summary estimates and 95% CIs with inverse variance weighted fixed-effects meta-analyses because heterogeneity was low²⁴ and random-effects meta-analysis might apply inappropriately large weights to smaller studies.²⁵ We characterised heterogeneity with the *I*² statistic. We used Cochran’s *Q* statistic to test for subgroup interactions and χ^2 tests to test for trend in analyses stratified by baseline systolic blood pressure. All p values were calculated from two-tailed tests of statistical significance with a type I error rate of 5%.

We did analyses to establish, first, the effect of a 10 mm Hg blood pressure reduction on the relative risk of major cardiovascular disease, coronary heart disease, stroke, heart failure, renal failure, and all-cause mortality; second, the effect of a 10 mm Hg blood pressure reduction at different baseline blood pressure levels by stratification of trials into five strata of reported mean baseline systolic blood pressure at the trial level (<130, 130–139, 140–149, 150–159, and \geq 160 mm Hg); third, the effects of a 10 mm Hg blood pressure reduction on the relative risk of major cardiovascular disease, coronary heart disease, stroke, heart failure, renal failure, and all-cause mortality in the presence of baseline comorbidities (cardiovascular disease, coronary heart disease, cerebrovascular accident, type 2 diabetes, heart failure, and chronic kidney disease) by investigation of possible interactions in treatment effect by these comorbidities; and fourth, the effects of different classes of blood pressure lowering drugs.

For the first three objectives, we standardised the analyses to a 10 mm Hg reduction in systolic blood pressure because we were interested in the proportional effects of lowering systolic blood pressure by 10 mm Hg and because the trials varied in the relative intensity of blood pressure lowering achieved due to differences in strategies and drugs. We standardised the analyses by multiplying the log of the summary statistic of each trial (and its standard error) by 10/d, where d was the average systolic blood pressure reduction in that trial.

For example, if the logHR was -0.2 and the systolic blood pressure reduction was 4 mm Hg, the standardised logHR would be $-0.2 \times (10/4) = -0.5$. We examined non-standardised effects of blood pressure lowering by baseline systolic blood pressure in sensitivity analyses.

We did meta-regression to assess the validity of the assumption that reductions in RR would be proportional to the achieved blood pressure reduction. The percentage reductions in proportional risk of major cardiovascular disease events, coronary heart disease, stroke, heart failure, renal failure, and all-cause mortality were regressed against the difference in mean achieved systolic blood pressure between the intervention and control groups.

Trials of blood pressure lowering drugs versus placebo or higher versus lower blood pressure targets were combined for the purposes of the first three objectives. For the six trials with three arms, including two active groups and a placebo group^{10,26–31} and the one trial that had four arms including a placebo group,³² we combined the active groups for the blood pressure lowering analysis by combining the events and taking a weighted average of baseline blood pressure, achieved blood pressure, and blood pressure reduction.

For the fourth objective, the comparison of drug classes, we did not standardise the analyses in order to account for variations in blood pressure lowering efficacy, tolerability, or non-blood pressure-mediated effects of the different drug classes. In these analyses, we examined possible differences in the effects of blood pressure lowering by drug class by comparing trials that tested a specific class of drug (angiotensin-converting enzyme [ACE] inhibitors, angiotensin receptor blockers [ARB], β blockers, diuretics, and calcium channel blockers [CCB]) against all other classes that it has been compared with. Because the pooled comparators might vary for each class (for example, β blockers might have mostly been tested against CCBs, whereas diuretics might have mostly been tested against ACE inhibitors), we also did an analysis in which we compared each individual drug class to all of the individual classes it has been tested against.




Figure 4: Standardised effects of a 10 mm Hg reduction in systolic blood pressure stratified by blood pressure. Blood pressure strata are baseline blood pressure values, not achieved blood pressure after treatment. RR=relative risk.

We excluded trials that were done in populations with heart failure or left ventricular systolic dysfunction from all of the main standardised analyses, because the effects of antihypertensive use on achieved blood pressure in heart failure have been reported to vary substantially with baseline blood pressure within the same trial population (thus rendering mean achieved

blood pressure less meaningful).³³⁻³⁵ Furthermore, only a few heart failure trials have actually reported mean achieved blood pressure by treatment allocation. However, we did complementary sensitivity analyses that included heart failure trials (both standardised and non-standardised) with the available information from published reports.




Figure 5: Standardised effects of a 10 mm Hg reduction in systolic blood pressure stratified by history of cardiovascular disease
 Data are stratified by subgroups in which all (cardiovascular disease) or none (no cardiovascular disease) of the participants had a history of cardiovascular disease at baseline. A cardiovascular disease subgroup is not shown for renal failure because no trial that reported renal failure as an outcome reported an analysis stratified by the presence of cardiovascular disease. RR=relative risk.

We did all analyses with Stata version 13.1 and R version 3.2.0.

Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. KR and DE had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Results

In total, we screened 11428 abstracts, of which 350 were eligible for full-text review (figure 1). Of the 134 randomised controlled trials identified, 123 trials with 613815 participants were eligible for inclusion in the meta-analysis. 92 studies were deemed to be trials of blood pressure lowering because they compared either blood pressure lowering drugs to placebo (78 trials) or different

blood pressure lowering targets (14 trials; appendix pp 1–6, 10). 43 trials compared different drug classes and were included in the drug comparison analysis (appendix pp 1–10). 12 trials fell into both categories, with five trials assessing different blood pressure lowering targets and drug classes^{36–42} and seven trials comparing blood pressure lowering drugs to placebo and different drug classes.^{26,28–30,32,43,44} Ten studies were judged to be of unclear risk of bias and 113 were deemed to be at low risk of bias.^{45–55} Heterogeneity for outcomes was low to moderate; the *I*² statistic for heterogeneity for major cardiovascular disease events was 41%, for coronary heart disease 25%, for stroke 26%, for heart failure 37%, for renal failure 28%, and for all-cause mortality 35%. The appendix shows details of the methods of blood pressure measurement for the included trials (appendix pp 11–14).

Meta-regression analyses showed relative risk reductions for major cardiovascular disease events




Figure 6: Non-standardised effects of reductions in systolic blood pressure stratified by class of blood pressure lowering drug
ACE=angiotensin-converting enzyme. ARB=angiotensin receptor blockers. CCB=calcium channel blockers. RR=relative risk.

($p < 0.0001$), stroke ($p < 0.0001$), heart failure ($p < 0.0001$), and all-cause mortality ($p = 0.014$) to be proportional to the magnitude of the blood pressure reduction achieved (figure 2, appendix pp 15–16). The meta-regression results were not significant for coronary heart disease ($p = 0.058$) or renal failure ($p = 0.09$; appendix pp 16–17).

Every 10 mm Hg systolic blood pressure reduction significantly reduced the risk of major cardiovascular disease events (RR 0.80, 95% CI 0.77–0.83), coronary heart disease (0.83, 0.78–0.88), stroke (0.73, 0.68–0.77), heart failure (0.72, 0.67–0.78), and all-cause mortality (0.87, 0.84–0.91; figure 3, appendix pp 18–23). The proportional reductions per 10 mm Hg systolic blood pressure reduction were greater for stroke and heart failure than for coronary heart disease. Estimates in non-

standardised analyses were consistent with standardised estimates (appendix p 24).

When we stratified trials by mean baseline systolic blood pressure and compared the effects of a 10 mm Hg reduction in systolic blood pressure between strata, we detected no significant trends for any outcomes ($p_{\text{trend}} > 0.05$; figure 4). Estimates were similar for non-standardised analysis (appendix p 25).

No evidence of different proportional effects of blood pressure lowering existed when we stratified trials by baseline cardiovascular disease (all $p_{\text{interaction}} > 0.05$; figure 5) and, in view of the multiplicity of tests done (which increases the likelihood of observing a chance finding when considering all tests), no strong evidence of differences existed when trials were stratified by

baseline coronary heart disease (all $p_{\text{interaction}} > 0.03$; appendix p 26). Although there was some evidence to suggest that the effect of blood pressure lowering on stroke was modified by presence of baseline coronary heart disease ($p=0.03$), we cannot rule out that this weak evidence is entirely due to chance given the multiplicity of testing. However, the proportional reduction in stroke risk seemed to be larger in populations without a history of cerebrovascular disease than in populations with such a history ($p_{\text{interaction}}=0.0028$; appendix p 27). When we stratified trials by baseline diabetes, we detected a significant interaction ($p=0.0006$) for major cardiovascular disease events, with significantly larger risk reductions for populations without diabetes (RR 0.75, 95% CI 0.70–0.80) than in populations with diabetes (0.88, 0.82–0.94; appendix p 28). Our subgroup analysis based on the presence of heart failure suggested that a 10 mm Hg blood pressure reduction might increase the risk of renal failure in patients with heart failure (21.67, 3.75–125.21), but this result was based on just 84 renal failure events in two studies (appendix pp 29–30). When we stratified trials by baseline chronic kidney disease, a significant interaction existed for major cardiovascular disease events ($p_{\text{interaction}}=0.012$), with larger proportional risk reductions for the populations without chronic kidney disease (0.68, 0.62–0.75) than in the populations with chronic kidney disease (0.84, 0.73–0.96). A significant interaction also existed for heart failure events, with a large and statistically significant risk reduction of 52% (0.48, 0.38–0.62) for every 10 mm Hg systolic blood pressure reduction in the subgroup without chronic kidney disease compared with a non-significant reduction for the subgroup with chronic kidney disease (0.95, 0.70–1.29; $p_{\text{interaction}}=0.0008$; appendix p 31).

We examined the five classes of blood pressure lowering drugs in non-standardised analyses. The different drug classes were of largely similar effectiveness for prevention of the various outcomes (figure 6, appendix pp 32–37). However, β blockers were less efficacious than other medications for the prevention of major cardiovascular disease events (RR 1.17, 95% CI 1.11–1.24; figure 6, appendix p 32), stroke (1.24, 1.14–1.35; figure 6, appendix p 33), and renal failure (1.19, 1.05–1.34; figure 6, appendix p 34). Evidence also suggested that β blockers had inferior efficacy in the prevention of all-cause mortality (1.06, 1.01–1.12; figure 6, appendix p 35), although this difference was not significant. CCBs were superior to the other classes for stroke prevention (0.90, 0.85–0.95; figure 6, appendix p 33) but were inferior to the other classes for heart failure prevention (1.17, 1.11–1.24; figure 6, appendix p 37). Diuretics were superior to other classes for heart failure prevention (0.81, 0.75–0.88; figure 6, appendix p 37). Results were similar when heart failure trials were included in the analysis (appendix p 38).

Discussion

In this meta-analysis, blood pressure lowering treatment significantly reduced the risk of cardiovascular disease and death in various populations of patients. Overall, a 10 mm Hg reduction in systolic blood pressure reduced the risk of major cardiovascular disease events by 20%, coronary heart disease by 17%, stroke by 27%, heart failure by 28%, and all-cause mortality by 13%. The size of these proportional reductions was broadly consistent across several major high-risk groups of patients, suggesting that blood pressure lowering provides broadly generalisable benefits. In stratified analyses, we saw no strong evidence that proportional effects were diminished in trials that included people with lower baseline systolic blood pressure (<130 mm Hg), and major cardiovascular events were clearly reduced in high-risk patients with various baseline comorbidities. Both of these major findings—the efficacy of blood pressure lowering below 130 mm Hg and the similar proportional effects in high-risk populations—are consistent with and extend the findings of the SPRINT trial.¹² Collectively, these data suggest that revision is urgently needed to recent blood pressure lowering guidelines that have relaxed the blood pressure lowering thresholds.^{14,17,18}

Our finding of a lack of overall benefit of blood pressure lowering for renal failure events is consistent with those of a previous meta-analysis⁵⁶ that assessed the effects of intensive versus moderate blood pressure reduction on the risk of end-stage kidney disease. Blood pressure lowering seems to have multiple and sometimes opposing effects on renal outcomes: long-term blood pressure lowering reduces proteinuria and other indicators of structural damage, especially but not exclusively when achieved by renin-angiotensin-aldosterone (RAAS) inhibitors, but increases in acute kidney injury have also been reported.^{12,57} These effects might be because of the fact that renal failure is not a single disorder, but a group of diseases with different underlying pathological mechanisms, with both high and low blood pressure contributing to its clinical manifestation.

A key insight from our analysis is that the effects of blood pressure lowering were broadly similar by baseline comorbidity. The proportional reduction in major cardiovascular disease events from blood pressure lowering did not differ substantially with the presence or absence of previous cardiovascular disease events, coronary heart disease, or cerebrovascular disease at the time of trial inclusion. With such consistent relative effects, we expect that the absolute benefits of blood pressure lowering would be greatest among individuals at highest absolute risk of cardiovascular events.^{56,58} However, we did note some differences if diabetes or chronic kidney disease were present at baseline. The proportional reduction in major cardiovascular disease events seemed to be larger in trials done in people without diabetes or chronic kidney disease. The cause of these differing proportional reductions is unclear. One possibility is that

the trials done in people with diabetes or chronic kidney disease differed in their methodological characteristics, such as length of follow-up or use of dual renin-angiotensin system inhibition.⁵⁹ Another possibility is that the non-significant finding of the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial, which was weighted heavily in the diabetic subgroup analysis, might have been caused by chance.⁵⁷

Another key finding from our analysis was that, although different drug classes were of largely similar effectiveness for prevention of the vascular outcomes of interest, there was evidence of modest differential effects between drug classes. β blockers seemed to be inferior to other classes of blood pressure lowering drugs for the prevention of major cardiovascular disease events, stroke, renal failure, and all-cause mortality, whereas CCBs seemed to be inferior and diuretics superior for the prevention of heart failure to other classes.^{18,60,61} In line with previous research and current recommendations, CCBs seemed to be superior to other drug classes for stroke prevention.^{14,16,60} Although we detected small but significant differences between the effectiveness of particular classes for other outcomes, we cannot exclude the possibility that these effects might have arisen from differences in the control regimens, or by chance in view of the multiple comparisons done. Additionally, apparent differences between drug classes might have been modified by the concurrent use of multiple classes of blood pressure lowering drugs in many trials. For example, the ACE inhibitor and CCB combination has been investigated in a large trial⁶² and was reported to be more effective for prevention of major cardiovascular disease than was an ACE inhibitor and diuretic combination. Finally, certain drug classes might have particularly strong positive or negative effects in specific patient subgroups (eg, β blockers in the first years after a myocardial infarction¹⁶); an effect that we did not investigate in this review of the totality of evidence across different populations. Individual patient data meta-analyses of the type done by the BPLTTC might be able to address these questions in the future.

The broad consistency of the proportional effects of blood pressure lowering on cardiovascular outcomes across various baseline blood pressure levels and several disease categories will challenge the current guidelines on blood pressure and will support the case to shift their focus from rigid blood pressure targets to risk-based targets, even when starting systolic blood pressure is lower than 130 mm Hg.⁶³ Rather than a decision based on an arbitrary threshold for a single risk factor, this approach needs individualised assessment of the balance of absolute risks and benefits when physicians decide on the blood pressure level at which to start blood pressure lowering and the target blood pressure.⁶⁴ For this meta-analysis, adverse event data were too disparate and inconsistently reported to allow for formal analysis. However, our analysis did provide evidence against blood pressure lowering causing an increase in major cardiovascular events in patients with

previous disease and low baseline blood pressure, which was a concern raised by reports of J-curve associations.

Pooling data from 613 815 patients enrolled in 123 large-scale trials of blood pressure lowering, our study includes substantially more information than any previous meta-analyses that have addressed this question. Unlike in previous studies, we excluded no trials because of baseline comorbidities, thus allowing for greater generalisability of findings and an assessment of treatment effect stratified by the presence of various baseline comorbidities of interest.^{65,66}

With ageing populations, chronic kidney disease is becoming an increasingly prevalent and important public health problem, affecting 10–15% of the adult population.^{67–70} Our findings show significant relative risk reductions for both patients with and patients without chronic kidney disease. Although proportional risk reductions were smaller in patients with chronic kidney disease than in those without, in view of their higher absolute risks, substantial absolute benefits from blood pressure reduction can be achieved in this population.

A limitation of this meta-analysis is that we had no individual patient data, the use of which would have provided greater detail about the effects of blood pressure lowering at various levels of baseline systolic blood pressure. However, the eligible trials spanned a wide range of baseline blood pressure levels, allowing us to explore the effects of blood pressure lowering across a relatively wide range of blood pressures. A further limitation was that the eligible trials varied in several respects, including differences in trial populations, baseline comorbidities, and treatment regimens, and it is possible that methodological differences might have confounded the differences recorded across subgroups of trials. Although many studies have investigated differences in the clinical efficacy of various drug classes for cause-specific outcomes, relatively few studies have compared different drug combinations. Because most patients need combination therapy, identification of the optimum combinations of therapies might be more clinically relevant than investigating the effectiveness of single therapies. The fact that we had no individual patient data prevented such an analysis.

In conclusion, blood pressure lowering significantly reduces the risk of major cardiovascular disease events, coronary heart disease, stroke, heart failure, and all-cause mortality, with similar proportional reductions across various population subgroups, irrespective of starting blood pressure. Lowering of blood pressure into what has been regarded the normotensive range should therefore be routinely considered for the prevention of cardiovascular disease among those deemed to be of sufficient absolute risk.

Contributors

DE and KR had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. KR, CAE, and DE contributed to the study concept and design.

DE, CAE, SGA, and TC contributed to the acquisition of data. All authors analysed and interpreted the data. DE, CAE, and KR drafted the manuscript with critical revision for important intellectual content from all authors. DE and CAE did the statistical analysis. KR was the study supervisor.

Declaration of interests

AR reports grants from Servier and receives salary support in part from The George Institute for Global Health; George Health Enterprises, the social enterprise arm of The George Institute, has received investment for the development of fixed dose combination therapy containing statin, aspirin, and blood pressure lowering medications. JC has received research grants from Servier for the ADVANCE trial and ADVANCE-ON post-trial follow-up, administered through the University of Sydney, and honoraria for speaking about these studies at scientific meetings. All other authors declare no competing interests.

Acknowledgments

KR is supported by the National Institute of Health Research (NIHR) Oxford Biomedical Research Centre and an NIHR Career Development Fellowship. DE is supported by the Clarendon Fund and CAE is supported by the Rhodes Trust. The work of The George Institute is supported by the Oxford Martin School. SGA is an Academic Clinical Lecturer in Cardiology and is funded by the NIHR. We would like to thank Hisatomi Arima and Hiroaki Shimokawa for providing additional information relating to their studies, and Allan J Hsiao for support with translation.

References

- WHO. A global brief on hypertension. Geneva: World Health Organization, 2013.
- Lewington S, Clarke R, Qizilbash N, Peto R, Collins C. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet* 2002; **360**: 1903–13.
- Lawes CMM, Rodgers A, Bennett DA, et al. Blood pressure and cardiovascular disease in the Asia Pacific region. *J Hypertens* 2003; **21**: 707–16.
- Rapsomaniki E, Timmis A, George J, et al. Blood pressure and incidence of twelve cardiovascular diseases: lifetime risks, healthy life-years lost, and age-specific associations in 1.25 million people. *Lancet* 2014; **383**: 1899–911.
- Collins R, Peto R, MacMahon S, et al. Blood pressure, stroke, and coronary heart disease. Part 2. Short-term reductions in blood pressure: overview of randomised drug trials in their epidemiological context. *Lancet* 1990; **335**: 827–38.
- SHEP Cooperative Research Group. Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension. Final results of the Systolic Hypertension in the Elderly Program (SHEP). *JAMA* 1991; **265**: 3255–64.
- Staessen J, Fagard R, Thijs L, et al. Randomised double-blind comparison of placebo and active treatment for older patients with isolated systolic hypertension. *Lancet* 1997; **350**: 757–64.
- Beckett NS, Peters R, Fletcher AE, et al. Treatment of hypertension in patients 80 years of age or older. *N Engl J Med* 2008; **358**: 1887–98.
- Patel A, MacMahon S, Chalmers J, et al. Effects of a fixed combination of perindopril and indapamide on macrovascular and microvascular outcomes in patients with type 2 diabetes mellitus (the ADVANCE trial): a randomised controlled trial. *Lancet* 2007; **370**: 829–40.
- PROGRESS Collaborative Group. Randomised trial of a perindopril-based blood-pressure-lowering regimen among 6105 individuals with previous stroke or transient ischaemic attack. *Lancet* 2001; **358**: 1033–41.
- Yusuf S, Sleight P, Pogue J, Bosch J, Davies R, Dagenais G. Effects of an Angiotensin-Converting-Enzyme Inhibitor, Ramipril, on Cardiovascular Events in High-Risk Patients. *N Engl J Med* 2000; **342**: 145–53.
- Wright JTJ, Williamson JD, Whelton PK, et al. A randomized trial of intensive versus standard blood-pressure control. *N Engl J Med* 2015; **373**: 2103–16.
- Emdin C, Rahimi K, Neal B, Callender T, Perkovic V, Patel A. Blood pressure lowering in type 2 diabetes. *JAMA* 2015; **313**: 603–15.
- James P, Oparil S, Carter B, et al. 2014 Evidence-based guideline for the management of high blood pressure in adults report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA* 2014; **311**: 507–20.
- Wright JT Jr, Fine LJ, Lackland DT, Ogedegbe G, Dennison Himmelfarb CR. Evidence supporting a systolic blood pressure goal of less than 150 mm Hg in patients aged 60 years or older: the minority view. *Ann Intern Med* 2014; **160**: 499–503.
- Law MR, Morris JK, Wald NJ. Use of blood pressure lowering drugs in the prevention of cardiovascular disease: meta-analysis of 147 randomised trials in the context of expectations from prospective epidemiological studies. *BMJ* 2009; **338**: b1665.
- Mancia G, Fagard R, Narkiewicz K, Redon J, Zanchetti A, Bohm M. 2013 ESH/ESC guidelines for the management of arterial hypertension: the Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Eur Heart J* 2013; **34**: 2159–219.
- NICE. The clinical management of primary hypertension in adults: clinical guideline 127. London: National Institute for Health and Care Excellence, 2011.
- Liberati A, Altman D G, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions. *PLoS Med* 2009; **6**: e1000100.
- Kjaergard LL, Villumsen J, Gluud C. Reported methodologic quality and discrepancies between large and small randomized trials in meta-analyses. *Ann Intern Med* 2001; **135**: 982–89.
- World Health Organization—International Society of Hypertension Blood Pressure Lowering Treatment Trialists' Collaboration. Protocol for prospective collaborative overviews of major randomized trials of blood-pressure lowering treatments. *J Hypertens* 1998; **16**: 127–37.
- Annesi I, Moreau T, Lellouch J. Efficiency of the logistic regression and Cox proportional hazards models in longitudinal studies. *Stat Med* 1989; **8**: 1515–21.
- Higgins JP, Green S, eds. Cochrane Handbook for Systematic Reviews of Interventions. Chichester: John Wiley & Sons, 2008: i–xxi.
- Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003; **327**: 557–60.
- Baigent C, Peto R, Gray R, Parish S, Collins R. Large-scale randomized evidence: trials and meta-analyses of trials. In: Warrell D, Cox T, Firth J, eds. Oxford textbook of medicine, 5th edn. New York: Oxford University Press, 2010.
- Nissen SE, Tuzcu EM, Libby P, et al. Effect of antihypertensive agents on cardiovascular events in patients with coronary disease and normal blood pressure: the CAMELOT study: a randomized controlled trial. *JAMA* 2004; **292**: 2217–26.
- Hansson L, Zanchetti A, Carruthers SG, et al. Effects of intensive blood-pressure lowering and low-dose aspirin in patients with hypertension: principal results of the Hypertension Optimal Treatment (HOT) randomised trial. *Lancet* 1998; **351**: 1755–62.
- Lewis EJ, Hunsicker LG, Clarke WR, et al. Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med* 2001; **345**: 851–60.
- Medical Research Council Working Party. MRC trial of treatment of mild hypertension: principal results. *BMJ* 1985; **291**: 97–104.
- MRC Working Party. Medical Research Council trial of treatment of hypertension in older adults: principal results. *BMJ* 1992; **304**: 405–12.
- Chaturvedi V, Patel A, Chalmers J, et al. Effects of a perindopril-based blood pressure lowering regimen on cardiac outcomes among patients with cerebrovascular disease. *Eur Heart J* 2003; **24**: 475–84.
- Ruggenenti P, Fassi A, Ilieva AP, et al. Preventing microalbuminuria in type 2 diabetes. *N Engl J Med* 2004; **351**: 1941–51.
- Meredith PA, Östergren J, Anand I, et al. Clinical outcomes according to baseline blood pressure in patients with a low ejection fraction in the CHARM (Candesartan in Heart failure: Assessment of Reduction in Mortality and morbidity) Program. *J Am Coll Cardiol* 2008; **52**: 2000–07.
- Rouleau JL, Roecker EB, Tendra M, et al. Influence of pretreatment systolic blood pressure on the effect of carvedilol in patients with severe chronic heart failure: the carvedilol prospective randomized cumulative survival (COPERNICUS) study. *J Am Coll Cardiol* 2004; **43**: 1423–29.
- Anand IS, Tam SW, Rector TS, et al. Influence of blood pressure on the effectiveness of a fixed-dose combination of isosorbide dinitrate and hydralazine in the african-american heart failure trial. *J Am Coll Cardiol* 2007; **49**: 32–39.

- 36 Wright JT, Bakris G, Greene T, et al. Effect of blood pressure lowering and antihypertensive drug class on progression of hypertensive kidney disease: results from the AASK trial. *JAMA* 2002; **288**: 2421–31.
- 37 Estacio RO, Jeffers BW, Hiatt WR, Biggerstaff SL, Gifford N, Schrier RW. The effect of nisoldipine as compared with enalapril on cardiovascular outcomes in patients with non-insulin-dependent diabetes and hypertension. *N Engl J Med* 1998; **338**: 645–52.
- 38 Schrier RW, Estacio RO, Mehler PS, Hiatt WR. Appropriate blood pressure control in hypertensive and normotensive type 2 diabetes mellitus: a summary of the ABCD trial. *Nat Clin Pract Nephrol* 2007; **3**: 428–38.
- 39 Schrier RW, Estacio RO, Esler A, Mehler P. Effects of aggressive blood pressure control in normotensive type 2 diabetic patients on albuminuria, retinopathy and strokes. *Kidney Int* 2002; **61**: 1086–97.
- 40 Asayama K, Ohkubo T, Metoki H, et al. Cardiovascular outcomes in the first trial of antihypertensive therapy guided by self-measured home blood pressure. *Hypertens Res* 2012; **35**: 1102–10.
- 41 Holman R, Turner R, Stratton I, et al. Efficacy of atenolol and captopril in reducing risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 39. *BMJ* 1998; **317**: 713–20.
- 42 Turner R, Holman R, Stratton I, et al. Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. *BMJ* 1998; **317**: 703–13.
- 43 McMurray J, Solomon S, Pieper K, et al. The effect of valsartan, captopril, or both on atherosclerotic events after acute myocardial infarction: An analysis of the valsartan in acute myocardial infarction trial (VALIANT). *J Am Coll Cardiol* 2006; **47**: 726–33.
- 44 Yusuf S, Diener H, Sacco RL, et al. Telmisartan to prevent recurrent stroke and cardiovascular events. *N Engl J Med* 2008; **359**: 1225–37.
- 45 Hansson L, Lindholm LH, Niskanen L, et al. Effect of angiotensin-converting-enzyme inhibition compared with conventional therapy on cardiovascular morbidity and mortality in hypertension: the Captopril Prevention Project (CAPPP) randomised trial. *Lancet* 1999; **353**: 611–16.
- 46 Wilhelmsen L, Berglund G, Elmfeldt D, et al. Beta-blockers versus diuretics in hypertensive men: main results from the HAPPHY trial. *J Hypertens* 1987; **5**: 561–72.
- 47 Kanamasa K, Kimura A, Miyataka M, Takenaka T, Ishikawa K. Incidence of cancer in postmyocardial infarction patients treated with short-acting nifedipine and diltiazem. *Cancer* 1999; **85**: 1369–74.
- 48 Levine T, Bernink P, Caspi A, et al. Effect of mibefradil, a T-type calcium channel blocker, on morbidity and mortality in moderate to severe congestive heart failure: the MACH-1 study. *Circulation* 2000; **101**: 758–65.
- 49 National Intervention Cooperative Study in Elderly Hypertensives Study Group. Randomized double-blind comparison of a calcium antagonist and a diuretic in elderly hypertensives. *Hypertension* 1999; **34**: 1129–33.
- 50 Ogawa H, Kim-Mitsuyama S, Matsui K, Jinnouchi T, Jinnouchi H, Arakawa K. Angiotensin II receptor blocker-based therapy in Japanese elderly, high-risk, hypertensive patients. *Am J Med* 2012; **125**: 981–90.
- 51 Helgeland A. Treatment of mild hypertension: a five year controlled drug trial. The Oslo study. *Am J Med* 1980; **69**: 725–32.
- 52 Gong L, Zhang W, Zhu Y, et al. Shanghai trial of nifedipine in the elderly (STONE). *J Hypertens* 1996; **14**: 1237–45.
- 53 Liu L, Wang J, Gong L, Liu G, Staessen J. Comparison of active treatment and placebo in older Chinese patients with isolated systolic hypertension. *J Hypertens* 1998; **16**: 1823–29.
- 54 Wang JG, Staessen J, Gong L, Liu L. Chinese trial on isolated systolic hypertension in the elderly. Systolic Hypertension in China (Syst-China) Collaborative Group. *Arch Intern Med* 2000; **160**: 211–20.
- 55 Ogihara T, Matsuoka H, Rakugi H. Practitioner's trial on the efficacy of antihypertensive treatment in elderly patients with hypertension II (PATE-hypertension II study) in Japan. *Geriatr Gerontol Int* 2011; **11**: 414–21.
- 56 Xie X, Atkins E, Lv J, et al. Effects of intensive blood pressure lowering on cardiovascular and renal outcomes: a systematic review and meta-analysis. *Lancet* 2015; published online Nov 7. [http://dx.doi.org/10.1016/S0140-6736\(15\)00805-3](http://dx.doi.org/10.1016/S0140-6736(15)00805-3).
- 57 Cushman WC, Evans GW, Byington RP, et al. Effects of intensive blood-pressure control in type 2 diabetes mellitus. *N Engl J Med* 2010; **362**: 1575–85.
- 58 Di Angelantonio E, Kaptoge S, Wormser D, et al. Association of cardiometabolic multimorbidity with mortality. *JAMA* 2015; **314**: 52–60.
- 59 Parving H-H, Brenner BM, McMurray JJ, et al. Cardiorenal end points in a trial of aliskiren for type 2 diabetes. *N Engl J Med* 2012; **367**: 2204–13.
- 60 Neal B, MacMahon S, Chapman N. Effects of ACE inhibitors, calcium antagonists, and other blood-pressure-lowering drugs: results of prospectively designed overviews of randomised trials. *Lancet* 2000; **356**: 1955–64.
- 61 Thomopoulos C, Parati G, Zanchetti A. Effects of blood pressure lowering on outcome incidence in hypertension: 4. Effects of various classes of antihypertensive drugs—overview and meta-analyses. *J Hypertens* 2015; **33**: 195–211.
- 62 Jamerson K, Weber MA, Bakris GL, et al. Benazepril plus amlodipine or hydrochlorothiazide for hypertension in high-risk patients. *N Engl J Med* 2008; **359**: 2417–28.
- 63 Sundström J, Arima H, Jackson R, et al. Effects of blood pressure reduction in mild hypertension. *Ann Intern Med* 2015; **162**: 184–91.
- 64 Rahimi K, MacMahon S. Blood pressure management in the 21st century: maximizing gains and minimizing waste. *Circulation* 2013; **128**: 2283–85.
- 65 Thomopoulos C, Parati G, Zanchetti A. Effects of blood pressure lowering on outcome incidence in hypertension. 1. Overview, meta-analyses, and meta-regression analyses of randomized trials. *J Hypertens* 2014; **32**: 2285–95.
- 66 Turnbull F, Neal B, Algert C, Chalmers J, Woodward M, MacMahon S. Effects of different blood-pressure-lowering regimens on major cardiovascular events: results of prospectively-designed overviews of randomised trials. *Lancet* 2003; **362**: 1527–35.
- 67 Ninomiya T, Perkovic V, Turnbull F, et al. Blood pressure lowering and major cardiovascular events in people with and without chronic kidney disease: meta-analysis of randomised controlled trials. *BMJ* 2013; **347**: f5680.
- 68 Coresh J, Astor B, Greene T, Eknoyan G, Levey A. Prevalence of chronic kidney disease and decreased kidney function in the adult US population: Third National Health and Nutrition Examination Survey. *Am J Kidney Dis* 2003; **41**: 1–12.
- 69 Chadban S, Briganti E, Kerr P, et al. Prevalence of kidney damage in Australian adults: the AusDiab kidney study. *J Am Soc Nephrol* 2003; **14** (7 Suppl 2): S131–38.
- 70 Perkovic V, Cass A, Patel A, et al. High prevalence of chronic kidney disease in Thailand. *Kidney Int* 2008; **73**: 473–79.