

EDITORIAL

Vitamin D Supplementation and Increased Risk of Falling A Cautionary Tale of Vitamin Supplements Retold

Steven R. Cummings, MD; Douglas P. Kiel, MD, MPH; Dennis M. Black, PhD

The randomized clinical trial (RCT) by Bischoff-Ferrari et al¹ in this issue of *JAMA Internal Medicine* shows that vitamin D supplementation is associated with the risk of falls. Two “high” doses (60 000 IU of vitamin D₃ per month or 24 000 IU vitamin D₃ plus 300 mg of calcifediol per month) achieved a serum 25-hydroxyvitamin D (25[OH]D) level of 30 ng/mL in 80% of participants, a level that has been recommended as best for reducing the risk of fractures and for other health benefits (to convert 25[OH]D to nanomoles per liter, multiply by 2.496).^{2,3} However, compared with a dose of 24 000 IU of vitamin D₃ per month (equivalent to 800 IU per day), the higher doses had no effect on lower extremity physical performance and increased the risk of falls. A previous RCT⁴ in women of the same age showed that 500 000 IU of vitamin D per year achieved serum 25(OH)D levels of at least 30 ng/mL in most participants but significantly increased risk of falls by 15% and fractures by 26%.

A theoretical possibility has been raised that periodic administration of high doses of vitamin D accounts for the increased risk of falls and fractures.⁵ That hypothesis should be tested by placebo-controlled trials showing that a daily dose of 2000 IU, for example, reaching serum 25(OH)D levels of at least 30 ng/mL reduces the risk of falls and fractures.

The trial in this issue¹ had no placebo group and, therefore, could not test the effects of supplementation with 800 IU of vitamin D per day. However, a recent placebo-controlled trial by Uusi-Rasi et al⁶ found that an 800 IU per day supplement had no effect on physical function or risk of falls or injurious falls, whereas an exercise program reduced the risk of injurious falls by about half. Another trial found that 800 IU daily had no effect on lower extremity function or risk of falls in postmenopausal women 75 years or younger.⁷

It is uncertain whether any dose of vitamin D supplementation reduces the risk of falls or fractures in community-dwelling older adults. Previous meta-analyses of RCTs had differed about whether vitamin D supplements reduce the risk of falls⁸⁻¹¹ or fractures^{8,12-14} in community-dwelling elderly individuals. In contrast meta-analyses¹⁵⁻¹⁷ have shown that 800 IU of vitamin D and 1200 mg of calcium reduced the risk of hip fracture and mortality for patients dwelling in institutions. These patients should receive calcium and vitamin D supplements.

Clinicians should not recommend vitamin D supplements for other putative health benefits. There is no evidence from meta-analyses of RCTs that vitamin D supplementation reduces the risk of cardiovascular disease or cancer.^{13,18} In addition, a recent trial¹⁹ found that 1000 IU of vitamin D per day, with or without calcium, did not decrease the risk of colon cancer or recurrent adenomas in those with a history of colon adenomas.

The vitamin D story seems to be following the familiar pattern observed with antioxidant vitamins. Enthusiasm for the health benefits of vitamin supplements is coupled with the belief that “vitamins” are inherently safe and reinforced by observational studies showing, essentially, that healthy people have higher vitamin levels. Then RCTs and meta-analyses²⁰ proved that the supplements in fact increase mortality (β-carotene, vitamin E), or have no health benefits (vitamin A, vitamin C).

The strategy of supplementation with vitamin D to achieve serum levels of at least 30 ng/mL has not been established by RCTs to reduce the risk of falls and fractures. It may increase the risk of falling. Until that approach is supported by randomized trials with updated meta-analyses, it would be prudent to follow recommendations²¹ from the Institute of Medicine (IOM) that people 70 years or older have a total daily intake of 800 IU of vitamin D without routine measurement of serum 25(OH)D levels. It is prudent to get recommended intakes of vitamin D and other vitamins from a balanced diet with foods that naturally contain what is manufactured into supplements.

ARTICLE INFORMATION

Author Affiliations: California Pacific Medical Center Research Institute, San Francisco (Cummings); Department of Epidemiology and Biostatistics, University of California San Francisco (Cummings, Black); Institute for Aging Research, Hebrew SeniorLife, Department of Medicine, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, Massachusetts (Kiel).

Corresponding Author: Steven R. Cummings, MD, San Francisco Coordinating Center, Coordinating

Center, 185 Berry Street, Lobby 5, Suite 5700, San Francisco, CA 94107 (scummings@sccc-cpmc.net).

Published Online: January 4, 2016.
doi:10.1001/jamainternmed.2015.7568.

Conflict of Interest Disclosures: None reported.

REFERENCES

1. Bischoff-Ferrari HA, Dawson-Hughes B, Orav JE, et al. Monthly high-dose vitamin D treatment for the prevention of functional decline [published

online January 4, 2016]. *JAMA Intern Med*. doi:10.1001/jamainternmed.2015.7148.

2. Holick MF, Binkley NC, Bischoff-Ferrari HA, et al; Endocrine Society. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2011;96(7):1911-1930. doi:10.1210/jc.2011-0385.

3. Dawson-Hughes B, Mithal A, Bonjour JP, Boonen S, Burckhardt P, Fuleihan GE, et al IOF position statement: vitamin D recommendations for older

adults [published online April 4, 2010]. *Osteoporos Int*. doi:10.1007/s00198-010-1285-3.

4. Sanders MH, Newman AB, Haggerty CL, et al; Sleep Heart Health Study. Sleep and sleep-disordered breathing in adults with predominantly mild obstructive airway disease. *Am J Respir Crit Care Med*. 2003;167(1):7-14.

5. Dawson-Hughes B, Harris SS. High-dose vitamin D supplementation: too much of a good thing? *JAMA*. 2010;303(18):1861-1862. doi:10.1001/jama.2010.598.

6. Uusi-Rasi K, Patil R, Karinkanta S, et al. Exercise and vitamin D in fall prevention among older women: a randomized clinical trial. *JAMA Intern Med*. 2015;175(5):703-711. doi:10.1001/jamainternmed.2015.0225.

7. Hansen KE, Johnson RE, Chambers KR, et al. Treatment of vitamin d insufficiency in postmenopausal women: a randomized clinical trial. *JAMA Intern Med*. 2015;175(10):1612-1621. doi:10.1001/jamainternmed.2015.3874.

8. Bolland MJ, Grey A, Reid IR. Differences in overlapping meta-analyses of vitamin D supplements and falls. *J Clin Endocrinol Metab*. 2014;99(11):4265-4272. doi:10.1210/jc.2014-2562.

9. Bolland MJ, Grey A, Gamble GD, Reid IR. Vitamin D supplementation and falls: a trial sequential meta-analysis. *Lancet Diabetes Endocrinol*. 2014;2(7):573-580. doi:10.1016/s2213-8587(14)70068-3.

10. Murad MH, Elamin KB, Abu Elnour NO, et al. Clinical review: the effect of vitamin D on falls: a systematic review and meta-analysis. *J Clin Endocrinol Metab*. 2011;96(10):2997-3006. doi:10.1210/jc.2011-1193.

11. Bischoff-Ferrari HA, Dawson-Hughes B, Willett WC, et al. Effect of vitamin D on falls: a meta-analysis. *JAMA*. 2004;291(16):1999-2006.

12. Bolland MJ, Grey A. A case study of discordant overlapping meta-analyses: vitamin d supplements and fracture. *PLoS One*. 2014;9(12):e115934. doi:10.1371/journal.pone.0115934.

13. Bolland MJ, Grey A, Gamble GD, Reid IR. The effect of vitamin D supplementation on skeletal, vascular, or cancer outcomes: a trial sequential meta-analysis. *Lancet Diabetes Endocrinol*. 2014;2(4):307-320. doi:10.1016/s2213-8587(13)70212-2.

14. DIPART (Vitamin D Individual Patient Analysis of Randomized Trials) Group. Patient level pooled analysis of 68 500 patients from seven major vitamin D fracture trials in US and Europe. *BMJ*. 2010;340:b5463.

15. Chapuy MC, Arlot ME, Duboeuf F, et al. Vitamin D3 and calcium to prevent hip fractures in the elderly women. *N Engl J Med*. 1992;327(23):1637-1642.

16. Tang BM, Eslick GD, Nowson C, Smith C, Bensoussan A. Use of calcium or calcium in combination with vitamin D supplementation to

prevent fractures and bone loss in people aged 50 years and older: a meta-analysis. *Lancet*. 2007;370(9588):657-666.

17. LeBlanc ES, Zakher B, Daeges M, Pappas M, Chou R. Screening for vitamin D deficiency: a systematic review for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2015;162(2):109-122. doi:10.7326/m14-1659.

18. Chung M, Lee J, Terasawa T, Lau J, Trikalinos TA. Vitamin D with or without calcium supplementation for prevention of cancer and fractures: an updated meta-analysis for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2011;155(12):827-838. doi:10.1059/0003-4819-155-12-201112200-00005.

19. Baron JA, Barry EL, Mott LA, et al. A trial of calcium and vitamin D for the prevention of colorectal adenomas. *N Engl J Med*. 2015;373(16):1519-1530. doi:10.1056/NEJMoa1500409.

20. Bjelakovic G, Nikolova D, Glud C. Meta-regression analyses, meta-analyses, and trial sequential analyses of the effects of supplementation with beta-carotene, vitamin A, and vitamin E singly or in different combinations on all-cause mortality: do we have evidence for lack of harm? *PLoS One*. 2013;8(9):e74558. doi:10.1371/journal.pone.0074558.

21. Institute of Medicine. *Dietary Reference Intakes for Calcium and Vitamin D*. Washington, DC: National Academies Press; 2011.